

‘NextGen’ Projections for the Western Tropical Pacific: Current and Future Climate for Cook Islands

Technical Report



October 2021



CSIRO and SPREP (2021). 'NextGen' Projections for the Western Tropical Pacific: Current and Future Climate for Cook Islands. Final report to the Australia-Pacific Climate Partnership for the Next Generation Climate Projections for the Western Tropical Pacific project. Commonwealth Scientific and Industrial Research Organisation (CSIRO) and Secretariat of the Pacific Regional Environment Programme (SPREP), CSIRO Technical Report, Melbourne, Australia. <https://doi.org/10.25919/24z1-tv40>

Acknowledgements

This report has been prepared for the Australia-Pacific Climate Partnership (APCP) as part of the APCP-funded project entitled Next Generation Climate Projections for the Western Tropical Pacific. The authors are as follows:

Michael Grose (CSIRO)
Savin Chand (Federation University)
Anil Deo (Federation University)
Leanne Webb (CSIRO)
Kim Nguyen (CSIRO)
Kathy McInnes (CSIRO)
Xuebin Zhang (CSIRO)
Julian O'Grady (CSIRO)
Rebecca Gregory (CSIRO)

The project team would like to acknowledge the technical support of the Cook Islands Meteorological Service for preparation of this report. The project team would also like to acknowledge Dr James Renwick (Victoria University of Wellington, New Zealand), Dr Yuji Masutomi (National Institute for Environmental Studies, Japan) and Dr Christophe Menkes (French National Research Institute for Sustainable Development, New Caledonia) for kindly undertaking external peer review of the draft final report, and Dr Dewi Kirono, Mr Kevin Hennessy and Dr Geoff Gooley (CSIRO) for undertaking the internal (CSIRO) review of this report.

© Commonwealth Scientific and Industrial Research Organisation 2021. To the extent permitted by law, all rights are reserved and no part of this publication covered by copyright may be reproduced or copied in any form or by any means except with the written permission of CSIRO.


Important disclaimer

CSIRO advises that the information contained in this publication comprises general statements based on scientific research. The reader is advised and needs to be aware that such information may be incomplete or unable to be used in any specific situation. No reliance or actions must therefore be made on that information without seeking prior expert professional, scientific and technical advice. To the extent permitted by law, CSIRO (including its employees and consultants) excludes all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses and any other compensation, arising directly or indirectly from using this publication (in part or in whole) and any information or material contained in it.

CSIRO is committed to providing web accessible content wherever possible. If you are having difficulties with accessing this document please contact csiro.au/contact

Contents

Executive Summary	i
1. Introduction	1
2. Historical and projected temperature	3
Historical change and observed variability	3
Different baselines	4
Climate stripes	5
‘Tracking’ projections from PACCSAP	6
Near-term projected variability and change	6
Projected change – near term, medium term and long term	8
Step-like changes	9
3. Historical and projected rainfall	10
Observed variability and change	10
Near-term variability and change	11
Projected change – near and long term	12
4. Projections for global warming levels	13
Temperature change	16
Rainfall change	16
5. Tropical cyclones, extreme rainfall and sea level projections	19
Tropical cyclones and extreme rainfall	19
Sea level	20
6. Standardised scenario analysis	24
Sampling the range of emissions and socio-economic pathways	24
Sampling the range of climate responses	25
Climate change storylines	26
Cook Islands North	27
Cook Islands South	28
7. Assessing impacts using these approaches	30
8. Data and Methods	32
9. References	35



Executive Summary

This report presents information about average temperature and rainfall change in Cook Islands, including historical change, interpretation of climate projections, understanding projections as they relate to 'global warming levels', and a set of future climate scenarios for Cook Islands using storylines. It also gives a summary of important new projections information on tropical cyclones, extreme rainfall and sea level rise, and gives a preview of the emerging set of new generation climate modelling.

This is a technical report for the Australia-Pacific Climate Partnership (APCP), as part of the APCP-funded project entitled 'Next Generation Climate Projections for the Western Tropical Pacific'. Technical reports of this type have been prepared for 14 Pacific Island countries plus Timor-Leste, upon which a suite of more user-friendly (and where possible non-technical) knowledge products have also been developed for communication, capacity development and general outreach to target users in the region. These users variously include national/sub-national governments, national met services and sectors, regional organisations and sectors, regional/national universities, consultancies, international donors and other development partners to inform climate-related policy development and risk assessments, adaptation and disaster risk management planning and associated decision-making. These reports, along with other project collateral including case study reports and project publications etc are available online at www.rccap.org.

This report uses much of the same underlying datasets and information from Pacific-Australia Climate Change Science and Adaptation Planning (PACCSAP) reports (CSIRO, Bureau of Meteorology and SPREP, 2014) but presents the information in new and more salient ways, with more context and detail; thereby making the underpinning science more relevant to decision-makers for application at a sectoral level. This report describes changes in average climate, which is superimposed upon natural year-to-year climate variability, partly related to the El Niño Southern Oscillation. The key findings for Cook Islands are summarised as follows:

Analysis of observed and future temperature

- Warming of the climate is clear and ongoing. From the 1850-1900 period (chosen to represent 'pre-industrial' climate), Cook Islands likely experienced around 0.5°C warming up to the 1986-2005 baseline and, due to warm years since 2000, around 0.7°C warming up to the 2011-2020 baseline period. The estimate is uncertain due to poor data early in the record.
- Observed warming trends from the 1986-2005 baseline to today are 'tracking' within the range of 2030 temperature projections described in the PACCSAP reports, suggesting that the PACCSAP projections are a reliable guide to the ongoing temperature trend.
- There is a long-term warming trend, but over the next ten years there may be a slight cooling trend or a rapid warming trend due to natural climate variability offsetting or enhancing the long-term trend.
- The warming is similar for the next 20 years regardless of the greenhouse gas emissions pathway (around 1.2°C from 1850-1900, or 0.7°C from 1986-2005), but later in the century, there is a growing and notable difference: less change under a low emissions pathway (RCP2.6) and much higher change under a higher emissions pathway (RCP8.5). By 2050 it's around 0.7°C (RCP2.6) to 1.3°C (RCP8.5), and by 2070 it's 0.8°C (RCP2.6) to 2.0°C (RCP8.5) (but slightly less in Cook Islands South), relative to 1986-2005.
- Step-like changes in temperature through time have occurred in the past, including an apparent step-increase since 2000, and we should expect there to be step-like changes in future.

Analysis of observed and future rainfall

- Historical rainfall trends are unclear given poor data coverage and very high climate variability. Also, the projected direction and magnitude of rainfall change is less clear than for temperature.
- There is a range of possible future changes in annual and seasonal rainfall, from wetter through to drier, largely determined by how the South Pacific Convergence Zone (SPCZ) changes and for the North region, the strength of the 'enhanced equatorial response' of warming along the equator. To plan for these possibilities, it is useful to assess the impact of both a wetter and a drier future – these are described in the standard climate scenarios with storylines (section 6).

Projections of change for ‘global warming levels’

Projections for Cook Islands are presented for a range of ‘global warming levels’ (defined as the warming of the global average temperature since the pre-industrial era), including the 1.5°C and 2°C limits from the Paris Agreement. In summary:

- During 2011-2020, the world was around 1.1°C warmer than in 1850-1900 (over land-only by around 1.6°C). Global warming could reach 1.5°C from the late 2020s or early 2030s. Whether we reach global warming levels higher than 1.5°C later in the century depends on the emissions pathway – from a very low pathway where the 2°C limit may not be crossed, to a very high pathway where 4°C or more is possible late in the century.
- In Cook Islands, observed and projected warming suggests that 2°C global warming relates to 1.4 to 1.9°C in Cook Islands North and 1.1 to 1.7°C in Cook South, i.e. Cook Islands warms at slightly less than the global warming rate. However, this doesn’t mean impacts are lower than other nations – high temperature extremes emerge faster in the tropics (Frame et al. 2017) and Cook Islands is vulnerable to climate change in many ways, so risk assessment is needed.
- A range of possible changes in average rainfall magnitude and direction are possible under all global warming levels, but the range of possible change is larger under higher global warming levels compared to lower levels.

New research into extremes and sea level

Since PACCSAP reporting in 2014, some updated research and assessments have been undertaken by the wider research community, and important new developments are summarised here for Cook Islands:

- **Extreme rainfall and tropical cyclones:** new research and assessment of evidence suggests that for the southwestern Pacific as a whole, the total number of tropical cyclones may decrease over the century, but with an increase in the average intensity and an increase in the impacts of tropical cyclones through more intense rainfall and greater coastal inundation due to sea level rise.
- **Sea level rise:** research since 2014 that was assessed in the IPCC Special Report on Oceans and

Cryosphere in a Changing Climate (SROCC, IPCC 2019) indicates that under a high global emissions pathway, the Antarctic ice sheet may contribute to greater sea level rise this century than previously thought. This means a higher high range of projected change is estimated – for Cook Islands this means an updated projection by 2090 under very high emissions of 0.51 to 0.98 m consistent with SROCC. Results are not as different for lower emissions pathways.

Preview of emerging emissions pathways and climate modelling

A set of new standard global emissions pathways are now defined, and a new set of coordinated global climate modelling is nearly finalised, emerging in 2021-2022. Important features of these new developments include:

- **Emissions Pathways:** we currently use the Representative Concentration Pathways (RCPs), but soon most work will move to the new Shared Socio-economic Pathways (SSPs), which include detailed emissions pathways based on the latest research but also include a global socio-economic dimension.
- **New models (CMIP6):** many climate projections use the CMIP5 set of global climate models, and a new set of CMIP6 models is emerging in 2021 and 2022. The new models show improvements at simulating the climate of the western Pacific. This means the confidence in projections may be a little higher in some instances – however many issues remain and the projections still show a wide range of possible change, so we still need to take a scenario approach and consider different future possibilities when planning climate adaptation.
- **Climate sensitivity:** the new set of models has a wider range of ‘climate sensitivity’ (the response of the climate to changes in greenhouse gases) than the existing set, with some models showing very high values and high temperature projections, but others showing very low climate sensitivity and low temperature change. These models are being assessed, and we may need to use this set of models differently than CMIP5 to produce projections to account for this broader range.

Standardised scenarios with storylines

For Cook Islands, a set of four representative future climate scenarios for 2050 is presented as a simple and effective means to represent the range of possibilities. The four scenarios are shown below in a 2x2 matrix with two dimensions:

- **Rows:** the range of possible emissions pathways the world follows, noting the likely socio-economic pathway. From a very low emissions pathway with a socio-economic pathway of ‘sustainability’ (top row) to a very high emissions pathway with a socio-economic pathway of ‘fossil-fuelled development’ (bottom row)
- **Columns:** two different but plausible changes in the regional climate for each of the emissions pathways, from warmer and drier (left column) to hotter and wetter (right column), determined by the dominant physical climate ‘storyline’ the region follows – for Cook Islands North this is if the South Pacific Convergence Zone (SPCZ) moves south compared to if it moves north and how strong the enhanced equatorial response is. For the South this relates to just changes in the SPCZ.

These four scenarios can be used as a simple starting point for considering the possible future climate in any risk assessment. Studies can compare the top and bottom rows (high and low emissions) indicating the benefits of following a low emissions pathway rather than a high pathway. Studies can also compare the climate scenario columns showing quite different possible outcomes for each emissions pathway.

The projected changes reported here can be applied to daily and monthly climate datasets representing the local conditions to examine the impact of climate change. Various analyses are possible, such as:

1. Subtracting past changes to examine change at different periods in the past
2. Adding on projected future changes to examine future scenarios, such as:
 - The complete range of plausible temperature change to 2050
 - Changes in temperature and rainfall for the four standardised scenarios
 - Changes in temperature and rainfall when the world reaches 2°C global warming since the pre-industrial era

Standardised scenarios for the Cook Islands for the period 2040-2059 relative to 1986-2005 for low and high emission pathways and two climate change scenarios defined by the physical change ‘storyline’.

	Scenario 1 SPCZ moves south Lower equatorial warming	Scenario 2 SPCZ moves north Higher equatorial warming
Low emissions (RCP2.6)	<p>Warmer & drier</p> <ul style="list-style-type: none"> • Annual temperature: +0.5°C • Annual rainfall: -10 to -15% • More heatwaves • Less humidity • More solar radiation • Heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 16-29 cm 	<p>Much warmer & wetter</p> <ul style="list-style-type: none"> • Annual temperature: +1.0°C • Annual rainfall: +10 to +15% • More heatwaves • More humidity • Less solar radiation • Much heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 16-29 cm
High emissions (RCP8.5)	<p>Much warmer & drier</p> <ul style="list-style-type: none"> • Annual temperature: +1.0°C • Annual rainfall: -15 to -20% • More heatwaves • Less humidity • More solar radiation • Heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 19-36 cm 	<p>Hotter & much wetter</p> <ul style="list-style-type: none"> • Annual temperature: +1.7°C • Annual rainfall: +15% • Many more heatwaves • More humidity • Less solar radiation • Much heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 19-36 cm

1 Introduction

This report presents new analyses and explanatory descriptions of historical and projected future temperature and rainfall to improve the interpretation and translation of climate change information in policy and sectoral planning. The focus is on delivering climate projections for Cook Islands and Pacific Island Countries (PICs) in the form of more contextualised science-based services using presentation styles tailored specifically for policy and sectoral user needs. This aims to facilitate appropriate and more extensive use of science-based evidence for on-ground applications.

This work has been undertaken as part of the project entitled 'Next Generation Climate Projections for the Western Tropical Pacific' funded by the Australian Government through the Australia-Pacific Climate Partnership (APCP) and delivered by the CSIRO Climate Science Centre (CSC) in partnership with the SPREP Pacific Met Desk Partnership.

The research builds upon existing climate projections, data and information developed previously through the Pacific Climate Change Science Program (PCCSP; Australian Bureau of Meteorology and CSIRO 2011) and the Pacific-Australia Climate Change Science and Adaptation Planning (PACCSAP) program (CSIRO, Australian Bureau of Meteorology and SPREP 2014). This report describes changes in average climate, which is superimposed upon natural year-to-year climate variability, partly related to the El Niño Southern Oscillation.

The work addresses new needs from climate projections, and it uses new research findings from updated observed datasets and draws on lessons from a series of sectoral engagements within Cook Islands and other PICs. Initial plans and results for the updated projections were presented at a number of forums, including a Pacific-regional workshop in Samoa, and national and sub-national workshops in Cook Islands. These engagements identified new and emerging data and information needs for informing climate risk and associated decision-making at the national level, and this was used to guide final analyses and production of this report.

New analyses of the average annual temperature and seasonal rainfall include:

- **Historical variability and change:** as far back as the 'pre-industrial' baseline 1850-1900
- **Change between different historical periods (baselines)**
- **Climate stripes:** visualisation of change from blue years (cooler than average) to red years (warmer than average)
- **'Tracking' projections:** assessing projections from a 20-year period centred on 1995 (1986-2005) to a 20-year period centred on 2030 (2020-2039) presented in PACCSAP
- **Near-term projected variability and change:** considering climate over the next 10 years
- **Projected change:** projections from different baselines and under different emissions pathways
- **Step-like projected changes:** assessing the possibility of step-like changes in the past and future
- **An example of applying these changes to regional analysis**

Other highlights:

- Projected changes are also presented for different 'global warming levels' from pre-industrial, including what the 1.5 and 2°C limits from the Paris Agreement mean for Cook Islands.
- Standardised future climate scenarios for 2050 with attached 'storylines' (descriptions of the factors that determine the future climate of the region).

The report also gives a brief overview of new developments in tropical cyclone (TC), extreme rainfall and sea level projections and gives an initial assessment of the new global emissions scenarios and climate modelling currently emerging.

Cook Islands

The Cook Islands are an archipelago of 15 major islands between 14 °S to 22 °S Latitude and 170 °W to 155 °W Longitude in the western Pacific Ocean (Figure 1.1). The northern group of islands lies to the north of the South Pacific Convergence Zone (SPCZ), the southern group lies to the south of the SPCZ. There are several key weather stations with long-term climate records in Cook Islands, including Rarotonga and Penrhyn and analysis of gridded observations and models reported here are average values calculated over the entire exclusive economic zone of Cook Islands, in two sub-regions including land and ocean (Figure 1.1, right). Both sub-regions of the Cook Islands have a wet season from November to April and dry season from May to October (Figure 1.2).



Figure 1.1. Map of Cook Islands (left) and the mask area (right) for calculating areal averages from gridded observational data and climate models' outputs.




Figure 1.2. Mean annual cycle of rainfall and temperature in Cook Islands in 1981-2010 (rainfall uses ERA5, temperature uses HadCRUT5, see Methods in Section 8).

2 Historical and projected temperature

For Cook Islands North and South, a projected average warming of around 0.7°C (0.4 to 1.0°C uncertainty range) from the 1986-2005 period out to 2020-2039 under all emissions pathways was reported (Australian Bureau of Meteorology and CSIRO, 2014). By 2090, the North is projected to warm by 2.0 to 3.8°C for very high emissions (RCP8.5) and 0.5 to 1.2°C for very low emissions (RCP2.6), similar in the South (1.7 to 4.2°C, and 0.2 to 1.1°C). What does this mean in the context of natural variability we have seen in the past, and how may this projection play out in the future?

Historical change and observed variability (Figure 2.1): average annual temperature has year-to-year variability, with cool years and warm years, but both sub-regions have showed a warming trend over the 1850-2020 period as a whole. There were periods of more rapid warming such as around 1980–2010 in the South, and less rapid warming or even cooling such as around 1950–1980 in the South. It appears likely that almost all years since 2000 are warmer than the pre-industrial climate average. Be aware that weather station data in the region are sparse before 1950 and particularly before 1920, so the data from these earlier periods are less reliable.




Figure 2.1. Average annual temperature of the Cook Islands North and South regions relative to 1850-1900 in five global datasets (see Methods, Section 8 for details).

Different baselines (Figure 2.2): The temperature of the Cook Islands region has risen, and at a faster rate in recent decades. This can be seen in increasingly warm temperatures through commonly used historical periods (or 'baselines').

This change between different baseline periods is important in three ways:

- It shows warming is underway and ongoing – we can't assume temperature is stable over time, and we must report any differences in context of the baseline used.
- The early pre-industrial baseline of 1850-1900 is now used widely to approximate a pre-industrial climate, so we must allow for warming that has already occurred since that time.
- Warming in Cook Islands has been less than the global average – from the 1850-1900 baseline up until the last 10 years, the world (land and ocean) warmed by around 1.1°C and land by around 1.6°C, but Cook Islands warmed by around 0.7°C.




Figure 2.2 Average annual temperature of Cook Islands North and South regions relative to 1850-1900 (°C; grey band indicates the range of five global temperature datasets) and the climate average for four different historical periods (baselines), as marked:

- WMO 1961-1990 is a common baseline used by the World Meteorological Organisation
- AR5 1986-2005 is the baseline used in the IPCC 5th assessment report (AR5), and PACCSAP
- AR6 1995-2014 is the new baseline that has been used in the IPCC 6th assessment report (AR6)
- 2011-2020 is a recent ten-year period

Climate stripes (Figure 2.3): The stripe pattern developed by Hawkins (2018) gives an indication of the variability, or ups and downs, in the temperature record. For Cook Islands, we see a clear change from more blue (cooler than average) to more red (warmer than average) years since 1850, with more red bars, especially since 2000.




Figure 2.3 Cook Islands North and South regions temperature relative to 1961-1990 through time presented as ‘climate stripes’ devised by Hawkins (2018): red = hotter, blue = cooler (using the Berkeley dataset).



© Salea Nihmei

‘Tracking’ projections from PACCSAP (Figure 2.4): temperature projections presented in the PACCSAP report are relative to the 1986-2005 historical period (or ‘baseline’), a period centred on 1995. We can ask the question “are the projections getting it right so far?” This graph shows that since 1995, the observed temperature change is within the range of what was projected by the climate models, suggesting the projections are a good guide to what’s happening.




Figure 2.4 Average annual temperature of the Cook Islands North and South regions relative to 1850-1900 (°C; grey band indicates range of five global temperature datasets), the reported range of projected change from PACCSAP is shown as a red wedge (the starting point allows for the difference between the five datasets) and the linear trend of smoothed observations for 1995-2020 plotted and extrapolated forward to 2030 (one dotted line for each dataset).

Near-term projected variability and change (Figure 2.5): as with the past climate, in future there always will be climate variability (ups and downs) at all time scales – including daily, monthly, yearly, 10-yearly and so on. We know that there is an ongoing warming trend, but that variability will occur as well. Two contrasting examples of how variability and the warming trend could unfold in Cook Islands North (results are similar for the South) are shown using two separate climate model simulations – these have a consistent underlying warming trend but a different sequence of climate variability.

The top plot shows a simulation where the linear trend in temperature in 2020-2030 is offset compared to the long-term trend, and in fact shows cooling. In contrast, the bottom panel shows a model simulation where the variability creates an enhanced temperature trend in 2020-2030 compared to the long-term trend. Note, these are not predictions of the next ten years, they are simulations shown to illustrate hypothetical cases. They emphasise the fact that we should always expect climate variability to enhance or offset the long-term trend.




Figure 2.5. Average annual temperature in Cook Islands North relative to 1850-1900 (°C) simulated in CMIP5 models, showing the range of all models (historic; grey band, future; blue band) and example model simulations (black line) with the linear trend for 2020-2030 marked (red line); top: an example with suppressed warming in 2020-2030 due to climate variability, bottom: an example where variability enhances the warming trend in 2020-2030.



© Salsesa Nihmei

Projected change – near term, medium term and long term (Figure 2.6): this graphic shows change under a very high emissions pathway in the pink shaded band (RCP8.5), and a very low emissions pathway in green (RCP2.6), with the model averages shown as thick lines. In the near term (2020-2039) the range of projected temperature change is similar for both emissions pathways, but in the medium term (2040-2059) the pathways begin to separate, and by the long term (2060-2079) the pathways give very different outcomes. By 2080 there is almost no overlap. By 2030, the warming is around 0.7°C. By 2050 it's around 0.7°C (RCP2.6) to 1.3°C (RCP8.5), and by 2070 it's 0.8°C (RCP2.6) to 2.0°C (RCP8.5) (but slightly less in Cook Islands South), relative to 1986-2005. Values are smaller from the recent baseline due to the historical warming already in place (see Table 4.1 for full details). Greenhouse gases have a long lifetime in the atmosphere, so early and sustained action taken to reduce emissions by the global community will reduce the climate change impacts experienced later.




Figure 2.6. Average annual temperature in Cook Islands North and South regions relative to 1850-1900 (°C) derived from observations (Berkeley shown; grey solid line) and simulated in CMIP5 models, showing the range of all models for the past period (grey band), the future under a very high emissions pathway (pink band) and a very low emissions pathway (green band). Thick horizontal lines show the mean of all models in 20-year periods of the baseline 1986-2005 (blue) and future 20-year periods centred on 2030, 2050 and 2070 (RCP8.5; red horizontal lines, RCP2.6; green horizontal lines).

Step-like changes (Figure 2.7): As well as year-to-year variability, and different 10-year trends (see above), there can be what look like ‘steps’ in the time series of mean annual temperature. As an example, in Cook Islands South (results are similar for the North), statistical tests detect two abrupt, step-like changes in the observed series of average temperature since 1850, making three ‘eras’ (shown as different colours in the plot). The most recent, warmest era is all years since 2000. Similarly, in model simulations we see apparent step-like changes and different eras in the past and future – with more steps and shorter eras during periods of more rapid background warming. Model simulations won’t predict the exact timing of these apparent steps but can indicate the kind of steps that may occur.

This illustrates that temperature change isn’t necessarily smooth, and this concept can help us better manage variability and change. We shouldn’t expect every year to be hotter than the last, and we should expect there to be not only variability year-to-year and between ten-year periods, but we should also expect what look like jumps or steps in the temperature record. This suggests that the climate system may have different stable states separated by abrupt changes or tipping points (Bathiany et al., 2018). The effect of these different apparent eras in the temperature series may be able to be seen in changes or impacts in the region.




Figure 2.7. Annual average temperature of Cook Islands South relative to the 1850-1900 (°C) baseline showing detected eras separated by breakpoints (see Data and Methods): top: in observations (Cowtan and Way 2014) showing three eras, early in the time series (blue), then ‘jump’ to a warmer period through mid to late 20th Century (orange), and the warmer 2000s (yellow); and bottom: an example model simulation, showing step-like changes throughout the past and future. Example model simulation is from ACCESS-1.0 under moderate emissions RCP4.5 (results are similar but with lower background warming in RCP2.6, and higher background warming in RCP8.5).

3 Historical and projected rainfall

Variability and percent change (%) in annual rainfall is examined here. PACCSAP reported rainfall projections for Cook Islands ranging from decrease to increase, with a multi-model average suggesting little change (near zero), except for the Cook Islands North region is projected to likely get drier in the May-October seas. Changes are potentially larger under higher emissions scenarios toward the end of the century. For example, the projected change for annual rainfall to 2030 ranges from -10 to +10% in all RCPs for North and South, but by 2090 the range is still within -10 to 10% under very low emissions (RCP2.6), and -15 to +25% under very high emissions (RCP8.5).

What does this mean in the context of natural variability we have seen in the past, and how will this play out in the future? Given the lack of a clear trend in observations, and the lack of clear projected direction of change, we don't show the 'tracking' or step-like change analyses, and just give a simple analysis of projections.

Observed variability and change (Figure 3.1): Annual total rainfall shows large year-to-year variability, partly related to the El Niño Southern Oscillation, and no significant trends since 1960 (Australian Bureau of Meteorology and CSIRO, 2014). Similarly, there are no significant trends in gridded rainfall datasets for the entire region of Cook Islands including surrounding oceans.

The average of all climate models considered (see Methods, Section 8) shows no significant change in annual rainfall for Cook Islands North and South between the pre-industrial baseline of 1850-1900 to the more recent baselines. Some individual models show increase, some decrease and the model average shows no change. This suggests there is no clear climate change signal and low model agreement in the historical period.




Figure 3.1 Area-average of Cook Islands North and South region annual rainfall (%) relative to the 1986-2005 period from gridded observations (ERA5 shown, results are similar from GPCP, CMAP and ERA5), the range of CMIP5 models (grey band). Coloured horizontal bars show the average of all models for the baseline periods described in the legend.

Near-term variability and change (Figure 3.2): large variability and no clear long-term trend means the 10-year trend in 2021-2030 could be for no change, decrease or increase. This is illustrated by the time series in two examples of model runs for Cook Islands South (results are similar for North).




Figure 3.2 Mean annual rainfall of the Cook Islands South region relative to 1986-2005 (%) from the group of CMIP5 models (grey band: historic, blue band: future), with an example model run shown in each panel (black line) highlighting the different possible trend in 2021-2030: (top) a model showing a decrease (red trend line in the top panel), and (bottom) a model showing an increase (red trend line in the bottom panel).



© Salea Nihmei

Projected change – near and long term (Figure 3.3): In contrast to temperature, there is a large range of possible change in mean rainfall from increase to decrease. While the multi-model mean is for little change for the near, medium and long term under both emissions pathways, this is just the mid-point between many models projecting increase and others projecting decrease. There is also very large variability in rainfall, making the definition of long-term trends unclear over shorter periods.




Figure 3.3. Average annual rainfall in the Cook Islands North and South regions region relative to 1850-1900 (%) in three gridded datasets using observations (coloured lines, as marked) and simulated in CMIP5 models, showing the range of all models for the past period (grey), the future under a very high emissions pathway (pink band) and a very low emissions pathway (green band). Thick lines show the mean of all models in 20-year periods: the baseline 1986-2005 (blue) and future 20-year periods centred on 2030, 2050 and 2070 (RCP8.5; red lines, RCP2.6; green lines).

4 Projections for global warming levels

It has been a common approach to describe future climate change in terms of what may occur by a defined future time-period, e.g. 'By 2030 warming for Cook Islands is projected to be around 0.7°C'. Recently there has been a growing focus on climate change according to different global warming levels – the amount that the world has warmed since the start of the industrial revolution. For example, the Paris Agreement aims to keep global warming to well below the level of 2°C since pre-industrial (1850-1900).

Global warming levels are measured as the global average surface temperature increase above the 1850–1900 baseline. The average global temperature for 2011–2020 is 1.1°C above this baseline, and a global warming of 1.5°C could occur in the window of the late 2020s to 2050 if warming continues at the current rate, and it will take dramatic emission reductions to avoid reaching it. Reaching higher global warming levels depends on the emissions pathway the world follows, the size of the global warming response to those emissions and the natural climate variability. This is shown by comparing the very low (top panel) and the very high (bottom panel) emissions pathways illustrated in Figures 4.1 and 4.2.




Figure 4.1 Global average surface temperature relative to 1850-1900 in Berkeley observations (grey line: annual value, red line: 41-year Lowess smoother), in the CMIP5 models for historical (grey band) and future under RCP2.6 (green band). The rough time window when global warming levels could be reached is shown by dark horizontal bars. Note models are calibrated relative to the 1986-2005 baseline and the observed change between 1850-1900 and 1986-2005 is added.




Figure 4.2 As for Figure 4.1 but showing a future under the very high emissions pathway (RCP8.5)

Because the projected warming rate varies by region, when the global average warming is at a defined level, some regions of the world will show warming lower than this average, some higher. The warming for Pacific Islands is projected to be lower than the global average, in contrast to the large continents and the Arctic that are projected to be much warmer. The maps below show the multi-model mean regional warming for the 1.5°C, 2°C, 3°C, and 4°C global warming levels (Figure 4.3).

Warming is less over the ocean regions than over land, so warming is lower over Cook Islands than over the large continents of Eurasia, North America and the Arctic (Figure 4.4). However, climate variability in the tropical Pacific is also lower than in many places, so in fact the 'signal to noise' ratio is similar to or higher than many other places. This means that unfamiliar climates, and higher temperature extremes can 'emerge' (become clear above natural variability) sooner (Frame et al. 2017; Hawkins et al. 2020). Also, it should be stressed the level of impact is determined by the climate hazards experienced, including extreme weather and climate events, but also on the current exposure and vulnerability of the country. In many ways Cook Islands is more vulnerable to the impacts of climate change than many other places.




Figure 4.3 The multi model average projection of temperature change across the world when the global average warming reaches 1.5, 2, 3 and 4°C above the 1850-1900 baseline.




Figure 4.4 As for the 2°C panel in Figure 4.3, but with a narrower colour scale that is centred on +2°C, with hotter than global average in red and cooler in purple.

Temperature change

The maps above show the multi-model average, but there is a range of regional temperature changes possible for each global warming level - suggested by different climate models. In a 2°C warmer world relative to the pre-industrial baseline, the Cook Islands are projected to be a little less than this global average: North 1.4 to 1.9°C, South 1.1 to 1.7°C warmer (Figure 4.5). Relative to the 1986-2005 baseline, the warming is about 0.7°C by 2030 for RCP2.6 and RCP8.5, by 2050 it's 0.7°C (RCP2.6) to 1.3°C (RCP8.5), and by 2070 it's 0.8°C (RCP2.6) to 2.0°C (RCP8.5) (but slightly less in Cook Islands South) (Table 4.1).




Figure 4.5. Change in the average annual temperature of the Cook Islands North and South regions at different global warming levels, from the 1850-1900 baseline, and from the more recent baseline of 1986-2005. The bars represent multi-model median and 10th-90th percentile range.

Rainfall change

We can examine rainfall projections for different global warming levels relative to the recent baseline period of 1986-2005. There is a range of future possibilities, with the median projection of little change in rainfall under any warming level for annual and 6-month season rainfall (Figure 4.6). The range of possible change is higher at higher warming levels. These results are consistent with the projections by timeframe and emissions pathway – where higher emissions pathways and further time periods bring higher global warming levels (Table 4.1).




Figure 4.6. Change in the average annual and 6-month seasonal rainfall in the Cook Islands North and South regions at different global warming levels relative to the 1986-2005 baseline. The bars represent multi-model median and 10th-90th percentile range.

Table 4.1a Projected changes in Cook Islands North average annual temperature and rainfall. Median changes are given, with the 10-90 percentile uncertainty range in brackets. Changes are for 20-year periods centred on 2030, 2050 and 2070, relative to 1986-2005, for low emissions (RCP2.6: green) and high emissions (RCP8.5: red). In 2030, changes are similar for low and high emissions. Projected changes in May-Oct rainfall and Nov-Apr rainfall are similar to those for annual rainfall.

	2030	2050	2070	1.5°C global warming	2°C global warming	3°C global warming	4°C global warming
Temperature from 1986-2005 (°C)	0.7 (0.4 to 1.0)	0.7 (0.5 to 1.2)	0.8 (0.4 to 1.2)	0.7 (0.4 to 0.9)	1.1 (0.8 to 1.3)	1.9 (1.5 to 2.1)	2.5 (2.0 to 3.0)
		1.3 (0.9 to 1.8)	2.0 (1.5 to 2.9)				
Annual rainfall from 1986-2005 (%)	0 (-8 to 9)	0 (-7 to 7)	-1 (-10 to 7)	1 (-4 to 7)	0 (-7 to 9)	2 (-11 to 12)	4 (-22 to 21)
		-1 (-14 to 14)	-2 (-7 to 9)				

Table 4.1b Projected changes in Cook Islands South average annual temperature and rainfall. Median changes are given, with the 10-90 percentile uncertainty range in brackets. Changes are for 20-year periods centred on 2030, 2050 and 2070, relative to 1986-2005, for low emissions (RCP2.6: green) and high emissions (RCP8.5: red). In 2030, changes are similar for low and high emissions. Projected changes in May-Oct rainfall and Nov-Apr rainfall are similar to those for annual rainfall.

	2030	2050	2070	1.5°C global warming	2°C global warming	3°C global warming	4°C global warming
Temperature from 1986-2005 (°C)	0.6 (0.3 to 1.0)	0.6 (0.3 to 0.9)	0.6 (0.3 to 1.1)	0.6 (0.3 to 0.7)	0.9 (0.6 to 1.1)	1.6 (1.3 to 2.0)	2.2 (1.8 to 2.6)
		1.2 (0.8 to 2.0)	1.8 (1.3 to 3.0)				
Annual rainfall from 1986-2005 (%)	1 (-10 to 10)	0 (-6 to 5)	1 (-3 to 6)	-1 (-5 to 7)	1 (-9 to 7)	3 (-8 to 11)	5 (-16 to 22)
		1 (-10 to 9)	2 (-11 to 20)				

5 Tropical cyclones, extreme rainfall and sea level projections

Tropical cyclones and extreme rainfall

Here we provide a brief overview of the latest research from this project and from around the world on tropical cyclone (TC) projections for the South Pacific region. Projections are given for a 2°C global warming since pre-industrial. As shown in the global warming levels section (Figure 4.1 and 4.2), this warming level could occur in around mid-century under RCP8.5, and there is approximately a 2/3 chance of staying below this target under RCP2.6 (IPCC, 2019). Changes may be larger under higher global warming levels.

Tropical cyclones can have dire socio-economic impacts on the South Pacific including Cook Islands, with damage to infrastructure and livelihoods arising from extreme winds and associated rainfall, coupled with destructive waves, storm surges and coastal flooding. Observations, theory and model experiments indicate, with increasing robustness, rising global-scale risks for some of these metrics that impact multiple regions. However, there is still a wide range in local- and regional-scale projections of TC activity (Figure 5.1). These large ranges are primarily due to challenges where climate models do not adequately resolve climatic processes and features affecting TC characteristics.




Figure 5.1 Southwest Pacific TC activity projection for a 2°C global warming. The bars on the left indicate a likely decrease in the total number of TCs, and a likely decrease but possible increase (but possible decrease) in the average intensity of TCs, and the very likely increase in the rainfall they bring. Shown are the median (blue line) and the 10th–90th-percentile ranges (blue bars). (Information used to construct this figure is derived from Knutson et al. 2020).

A recent study by Knutson et al. (2020) assesses the impact of a 2°C global warming on TC activity around the world. Below are some key messages from that study relevant to the broader southwest Pacific region; care must be exercised when interpreting these results for any local region in the Pacific, such as Cook Islands.

- High confidence that TC frequency will decrease over the coming century.
- Low confidence in the changes in frequency of category 4-5 TCs.
- High confidence that sea level rise will increase TC-related storm surge events.
- Medium to high confidence in an increase in TC rainfall rates.
- Medium to high confidence in the increase in average TC intensity.

The projected increase in average TC intensity, combined with sea level rise and increased rainfall rates would increase TC impacts. The reader is referred to Chand et al. (2020) for a review of tropical cyclones in the Pacific and implications of climate change.

Sea level

Research since 2014 was assessed in the IPCC (2019) Special Report on Oceans and Cryosphere in a Changing Climate (SROCC). Under a high global emissions pathway, the research suggested that Antarctic ice sheets may contribute to greater sea level rise this century than previously thought. Sea level projections that incorporate the higher Antarctic contribution have been evaluated for the Cook Islands and show a rise of between approximately 0.09 to 0.18 m by 2030 (very similar values for different RCPs and also similar to the PACCSAP projections), and an increase of 0.62 to 1.20 m by 2100 under RCP8.5 (Figure 5.2 and Table 5.1). Compared to the PACCSAP scenarios, the updated median value of sea level rise for RCP 8.5 in 2100 is 0.13 m higher and the upper end of the range is 0.16 m higher. This is because the enhanced Antarctic contribution is expected to be strongest for the highest emission scenario. For RCP 4.5 and 2.6 the range of projected change is similar but slightly narrower than PACCSAP projections. Interannual variability of sea level will lead to periods of lower and higher regional sea levels. In the past, this interannual variability (after removal of the seasonal signal) has been about 0.20 m (5–95% range; see dashed lines in Figure 5.2), and it is likely that a similar range will continue through the 21st century. The latest IPCC Sixth Assessment Report (IPCC 2021) emphasises that a ‘low likelihood high impact’ outcome of much higher sea level rise under the high emissions pathways can’t be ruled out.




Figure 5.2 Sea level rise projections for Cook Islands. The tide gauge record of relative sea level at Rarotonga is indicated in black, the satellite record in green and reconstructed sea level data is shown in purple, all are monthly means without seasonal cycles and referenced to mean sea level between 1986-2005. Multi-model-mean projections from 1995–2100 are given for the RCP8.5 (red solid line), RCP4.5 (cyan solid line) and RCP2.6 emissions scenarios (blue solid line), with the 5–95% uncertainty range for RCP8.5 and RCP2.6 shown by the red and blue shaded regions respectively. The dashed lines are an estimate of interannual variability in sea level (5–95% uncertainty range) and indicate that individual monthly averages of sea level can be above or below longer-term averages.

Table 5.1 Median sea level projections for Cook Islands with 5-95% range relative to 1986-2005 for RCPs 2.6, 4.5, and 8.5. Units are metres.

	RCP 2.6		RCP 4.5		RCP 8.5	
Year	Sea level rise		Sea level rise		Sea level rise	
2030	0.12	[0.09-0.17]	0.12	[0.09-0.16]	0.13	[0.09-0.18]
2040	0.17	[0.12-0.23]	0.17	[0.12-0.23]	0.19	[0.14-0.26]
2050	0.22	[0.16-0.29]	0.23	[0.16-0.31]	0.27	[0.19-0.36]
2060	0.27	[0.19-0.36]	0.29	[0.21-0.39]	0.35	[0.26-0.48]
2070	0.31	[0.23-0.42]	0.35	[0.26-0.48]	0.46	[0.33-0.63]
2080	0.36	[0.26-0.49]	0.42	[0.30-0.58]	0.58	[0.42-0.79]
2090	0.40	[0.29-0.55]	0.49	[0.35-0.67]	0.71	[0.51-0.98]
2100	0.45	[0.32-0.62]	0.56	[0.40-0.77]	0.86	[0.62-1.20]

Box 1. New emissions pathways

The international research community is set to move away from the Representative Concentration Pathways (RCPs) to the Shared Socio-economic Pathways (SSPs) in 2021. These new pathways are more relevant and up to date (e.g. they start in 2016, rather than in 2006), and they also include a description of the global social and economic factors that can produce each emissions pathway. There are five SSPs, but here we focus on the extreme ends of likely emissions – SSP1 and SSP5:

SSP1 – ‘Sustainability’, or ‘Taking the green road’ – higher challenges to mitigation but lower challenges to adaptation compared to some other SSPs. A world shifting to sustainable development with lower material growth, lower energy-intensity and lower resource-intensity, and along with this comes greater equity, inclusiveness, respect for environmental boundaries, education and health. Leads to the lower emissions pathways, so is most compatible with RCP2.6.

SSP5 – ‘Fossil-fuelled development’ or ‘Taking the highway’ – lower challenges to mitigation, higher challenges to adaptation compared to other SSPs. A world also investing in health, education and institutions, but one that relies on fossil fuels to drive rapid growth, technology and innovation to maintain energy-intensive and resource-intensive lifestyles. Leads to higher emissions pathways, so is most compatible with RCP8.5.

Others are:

- SSP2 – Middle of the road – medium challenges to mitigation and adaptation
- SSP3 – ‘Regional rivalry’ or ‘A Rocky Road’ – high challenges to mitigation and adaptation
- SSP4 – ‘Inequality’ or ‘A Road Divided’ – low challenges to mitigation, high challenges to adaptation


These are global scenarios, and further work is needed to see what they mean for a region or an individual country. For New Zealand and the Pacific, Frame et al. (2018) lay out some analysis of the SSPs and what they might mean for the region.

Box 2. New global climate models

A new group of global climate model simulations from around the world are being evaluated and will start to become the new international standard following the IPCC Sixth Assessment Report (2021). This means a shift from the Coupled Model Inter-comparison Project phase 5 (CMIP5) to the new phase, CMIP6.

The new models show incremental improvements in the simulation of the climate of the western Pacific, and there will be new insights and new opportunities to understand the future climate using these models. However, the fundamental nature of the models hasn't changed, and the western Pacific remains a challenging area to simulate climate variability and change, and confidence in regional climate projections will remain lower than for some other places.

As an example of the evaluation of models, this plot quantifies the well-known “cold-tongue bias” (the West Pacific warm pool is pinched in at the equator by a tongue of cold water) in the old and the new models:




Average sea surface temperature (SST) from 1950 to 2005 for June to November. Top panel shows observations (HadISST1.1) on a colour scale, and the thick black line shows 28.5°C. The other two maps show the CMIP5 and CMIP6 SST for the same period, and the thin grey lines show the location of 28.5°C for each individual model, and the thick black line shows the multi-model means. Taken from Grose et al. (2020).

Also, the projected changes in CMIP6 are generally similar to CMIP5 for many features and many climate variables. For example, annual rainfall change under a very high emissions pathway (RCP8.5 or SSP5-8.5) shows many of the same aspects – high model agreement for a strong rainfall increase at the equator, but less certain direction of change for the SPCZ region. There is a group of models with higher temperature projections in CMIP6 compared to CMIP5, and these models are currently being assessed.

In summary, there will be value and new insights from CMIP6 compared to CMIP5, but the nature of the model biases and the projections remain broadly similar to CMIP5.

Projected change to average annual rainfall between 1950-1999 and 2050-2099 under a very high emissions pathway (36 models for RCP8.5 in CMIP5; 29 models available for SSP5-85 in CMIP6), showing the multi-model mean rainfall and change with stippling (dots) indicating where less than two thirds of the models agree on the direction of change – there is high agreement on increase at the equator, but lower agreement elsewhere in both model groups – but perhaps more agreement in CMIP6 in some key regions.



© Saleisa Nihmei

6 Standardised scenario analysis

The future of the climate is determined by three main factors:

- 1) Ongoing, natural climate variability
- 2) Greenhouse gas and aerosol emissions pathways due to human activities and
- 3) The change in the climate that results from the emissions (the 'climate response')

A standardised future climate scenario analysis is a simple and useful way of understanding and quantifying potential ranges of climate change and associated hazards in terms of the emissions pathways and climate responses. This information can then be used in climate impact or risk assessments, in addition to the impact of ongoing climate variability (see, e.g. Sections 3 and 4).

Standardised future climate scenarios need to be both **representative** and **internally consistent**:

- **Representative** means covering or sampling the full range of plausible future climates including the range of plausible emissions pathways (RCPs) and the range of plausible climate responses to each emissions pathway – assessed using multiple lines of evidence including the results from different climate models.
- **Internally consistent** means that the changes in different climate variables (e.g. temperature and rainfall) make physical sense. When conducting risk assessments, a useful approach is to employ projections of different variables from a single climate model to ensure that projections are internally consistent. Mixing variables from different models (e.g. taking temperature from Model A while rainfall from Model B) into a single scenario may result in physically implausible combinations and is not recommended as best practice.

Here we describe a set of four representative, internally-consistent scenarios for the two regions of the Cook Islands for the 2050 period (2040–2059) relative to 1986–2005, covering the full plausible range of emissions pathways and climate responses (temperature and rainfall) as well as a description of the dominant physical 'storyline' of the climate change processes behind each case.


Please note that this is an illustration, it is not suitable for all purposes. A fully developed set of scenarios would need to be co-designed and co-produced between researchers and stakeholders in Cook Islands, for the relevant climate variables, timeframe, location and level of impact. The information would then be tailored for a risk assessment by those stakeholders. Please refer to the Pacific Guidelines (CSIRO and SPREP, 2017) on how to use and apply climate change information for impact assessment.

Sampling the range of emissions and socio-economic pathways

To sample each end of the range of global greenhouse gas emissions and global socio-economic changes (also called to 'bookend' or 'bracket' the range), we consider two cases:

- **High case** – the world is following a high emissions pathway (RCP8.5) on track for 3-4°C global warming by 2100 (or even more), likely under the global 'Fossil-fuelled development' shared socio-economic pathway SSP5 (see Box 1).
- **Low case** – the world is following a pathway to decarbonise the economy leading to net zero emissions by 2070 (RCP2.6), giving a two-thirds chance of staying below 2°C global warming by 2100, likely under the 'Sustainability' SSP1 (see Box 1).

The carbon dioxide concentrations given by different RCPs through the century are indicated by the different coloured lines shown in the below diagram. The circles show the two cases and give perspective on the pathways they represent in terms of carbon dioxide concentrations (a major greenhouse gas).



Sampling the range of climate responses

For the High case and the Low case emission pathways, the plausible range of change in temperature and rainfall can be assessed by looking at multiple well-performing climate models (Methods, Section 8) and each end of the change can be sampled (to ‘bookend’ or ‘bracket’ the range of possibilities here too).

The projections from all models can be plotted with temperature change on one axis and rainfall change on the other, producing a 2D shape or ‘uncertainty space’. This can be done for both the **Low Case** and **High Case** emission pathways (Figure 6.1).




Figure 6.1 Modelled change in annual temperature and rainfall between 1986-2005 and 2040-2059 from CMIP5 models (coloured shapes) in the Cook Islands region North at the left (N) and south at the right (S), showing the selected models that are representative of a warm dry future and a hot wet future for RCP2.6 (low case: green polygon) and RCP8.5 (high case: red polygon).

Bookending the temperature and rainfall projections requires two scenarios for each emission pathway: (1) warmer and drier, (2) hotter and wetter. Risk assessments often require information about other climate variables such as extreme daily temperature, extreme daily rainfall, extreme daily windspeed, drought, humidity, solar radiation and sea level rise. Internally consistent projections for some of these variables can be included under each scenario (Table 6.1). Changes in annual mean rainfall are typically correlated with changes in annual mean soil moisture, humidity and solar radiation. These are indicative scenarios for an initial scan of impacts, but detailed impact/risk assessments may need to consider a more comprehensive range of scenarios tailored for specific regions, sectors or systems.

For the **High Case** emissions pathway, the range of change in annual temperature and rainfall can be seen as time series in Figure 6.2. The separation between the time series grows with time.




Figure 6.2 Modelled annual temperature (°C) and rainfall (%) changes relative to 1986-2005 from CMIP5 models in the Cook Islands region North at the left (N) and south at the right (S), showing the selected models that are representative of a warm dry future (red) and a hot wet future (blue).

Table 6.1 Standardised scenarios for the Cook Islands for the period 2040-2059 relative to 1986-2005 for low and high emission pathways and two climate change scenarios defined by the physical change 'storyline'.

	Scenario 1* SPCZ moves south Lower equatorial warming	Scenario 2* SPCZ moves north Higher equatorial warming
Low emissions (RCP2.6)	Warmer & drier <ul style="list-style-type: none"> • Annual temperature: +0.5°C • Annual rainfall: -10 to -15% • More heatwaves • Less humidity • More solar radiation • Heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 16-29 cm 	Much warmer & wetter <ul style="list-style-type: none"> • Annual temperature: +1.0°C • Annual rainfall: +10 to +15% • More heatwaves • More humidity • Less solar radiation • Much heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 16-29 cm
High emissions (RCP8.5)	Much warmer & drier <ul style="list-style-type: none"> • Annual temperature: +1.0°C • Annual rainfall: -15 to -20% • More heatwaves • Less humidity • More solar radiation • Heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 19-36 cm 	Hotter & much wetter <ul style="list-style-type: none"> • Annual temperature: +1.7°C • Annual rainfall: +15% • Many more heatwaves • More humidity • Less solar radiation • Much heavier rainfall events • Greater tropical cyclone impacts • Sea level rise: 19-36 cm

* These are indicative scenarios for an initial scan of impacts, but detailed impact/risk assessments may need to consider a more comprehensive range of scenarios tailored for specific regions, sectors or systems.

Climate change storylines

We have more confidence in climate projections if there is a clear and convincing reason why the change would occur, drawing on multiple lines of evidence – this creates a climate ‘storyline’ of change backed up by evidence (Shepherd et al. 2018). If there is a range of possibilities that we can’t currently narrow down, then it is useful to identify the ‘storyline’ behind the each of the categories of possible change. In this way, we can plan for each of the possible future climates, while understanding what would cause them.

For temperature and rainfall change in both regions of the Cook Islands, there is a range of projected change, so we need to identify the plausible ‘storyline’ for each of the scenarios shown in the previous section. Due to its position on either side of the centre of the SPCZ, there is a different set of drivers for the North compared to the South, so a set of storylines is shown for each one.

Cook Islands North

The climate of Cook Islands North has a similar range of drivers as other countries of country sub-regions on or to the north of the centre of the SPCZ – so to look at this storyline, we group Samoa and Tuvalu together with Cook Islands North (see regions in Methods, Figure 8.2). The models that best represent the two storylines for the whole region, that also have published values of change in the SPCZ, are NorESM1-ME for warm and dry, and MPI-ESM-LR for hot and wet.

For temperature, the change for a given RCP and timeframe relates largely to the global climate sensitivity (how much the climate responds to a given increase in greenhouse gases). The warm and dry scenario has lower climate sensitivity and lower warming, while the hot and wet scenario has higher climate sensitivity and so higher warming. However, the change is also partly related to the change in rainfall (the shapes in Figure 6.1 above have the ‘warm dry’ to ‘hot wet’ diagonal orientation), so temperature change is partly related to changes in the major driver of rainfall too.

Rainfall change in this region is largely a function of changes in the location, angle and intensity of the dominant rainband, the SPCZ (Brown et al. 2013; Brown et al. 2020) as well as to the magnitude of the ‘enhanced equatorial response’ (how much the equator region becomes hotter and wetter, see Grose et al. 2014a). The countries in this region are on or at the northern edge of the SPCZ, so the movement of the SPCZ north or south will fundamentally change the rainfall they receive. The countries are also close to the equator, so a strong warming and rainfall increase at the equator will also affect the climate here.

Looking at the **High Case** in Figure 6.3, the warm dry scenario sees a southward movement and a weakening of the SPCZ, and a lower equatorial warming. In contrast, the hot wet scenario sees a northward movement and strengthening of the SPCZ, and a stronger equatorial response. The SPCZ changes in shape and extent, and together with the southerly movement, this means Cook Islands moves into a wetter zone.




Figure 6.3 The mean annual rainfall of the western tropical Pacific (bluer is wetter) in the 1986-2005 (past) and the 2040-2059 (future) periods as simulated in the models used to represent the warm and dry scenario (top) and the hot and wet scenario (bottom). The approximate position of the dominant rainband, the South Pacific Convergence Zone (SPCZ), is marked as a red line.

The result of these storylines can be seen as either a drying or a wetting across the wedge shaped region between the SPCZ and the equator (Figure 6.4). Both scenarios are plausible given our current understanding of processes driving climate change in the region. With further research, we may be able to have more certain projections of the SPCZ and the equator, reject non-plausible cases and present a narrower range of projected change. But for the moment we should consider both these scenarios, or anywhere in between, as possible. The scenarios are similar in nature but with smaller changes in the **Low Case**.




Figure 6.4 Map of projected change for mean annual temperature (left panels, °C) and rainfall (right panels, %) between 1986-2005 to 2040-2059 under RCP8.5 in the two scenarios for the **High Case**

Cook Islands South

The climate of Cook Islands has a similar range of climate drivers as other countries in the south SPCZ region – so to look at this storyline, we group Cook Islands South together with Fiji, Tonga, Niue and Vanuatu (see regions in Methods, Figure 8.2). The models that best represent the two storylines for the whole region, that also have published values of change in the SPCZ, are GISS-E2-R for warm and dry, and IPSL-CM5A-MR for hot and wet (similar models as used for Cook Islands, above).

For temperature, the change for a given RCP and timeframe relates largely to the global climate sensitivity (how much the climate responds to a given increase in greenhouse gases). The warm and dry scenario has lower climate sensitivity and lower warming, while the hot and wet scenario has higher climate sensitivity and so higher warming. However, the change is also related to the change in rainfall (the shapes in Figure 6.1 above have the ‘warm dry’ to ‘hot wet’ diagonal orientation), so temperature change is partly related to changes in the major driver of rainfall too.

Rainfall change in this region is largely a function of changes in the location, angle and intensity of the dominant rainband, the SPCZ (Brown et al. 2013; Brown et al. 2020). The countries in this region are at the southern edge of the SPCZ, so the movement of the SPCZ north or south will fundamentally change the rainfall they receive. This then has an impact on the warming they experience too.

Looking at the **High Case** in Figure 6.5, the warm dry scenario sees a northward movement of the SPCZ of 4° latitude over the century (Brown et al. 2013). Even though the SPCZ strengthens and extends, the northward movement causes the region of Cook Islands to move into a drier zone. In contrast, the hot wet scenario sees a southward movement of the SPCZ of 3° latitude over the century (Brown et al. 2013). The SPCZ changes in shape and extent, and together with the southerly movement, this means Cook Islands moves into a wetter zone.




Figure 6.5 The mean annual rainfall of the western tropical Pacific (bluer is wetter) in the 1986-2005 (past) and the 2040-2059 (future) periods as simulated in the models used to represent the warm and dry scenario (top) and the hot and wet scenario (bottom). The approximate position of the dominant rainband, the South Pacific Convergence Zone (SPCZ), is marked as a red line.

The result of these storylines can be seen as either a drying or a wetting across the broad south SPCZ region (Figure 6.6). Both scenarios are plausible given our current understanding of processes driving climate change in the region. With further research, we may be able to have more certain projections of the SPCZ, reject non-plausible cases and present a narrower range of projected change. But for the moment we should consider both these scenarios, or anywhere in between, as possible. The scenarios are similar in nature but with smaller changes in the Low case.




Figure 6.6 Map of projected change for mean annual temperature (left panels, °C) and rainfall (right panels, %) between 1986-2005 to 2040-2059 under RCP8.5 in the two scenarios for the High Case

7 Assessing impacts using these approaches

There are various historical and projected changes in this report, including:

- Historical changes between different baseline periods
- Future changes for different emissions scenarios and time frames
- Future changes for different global warming levels
- Future change in 2050 using standardised scenarios with storylines and representative climate models.

Here's how you can look in detail at what this means to the average climate of Cook Islands.

Observations at a weather station or as gridded data describe the local conditions today. If we want to see what a different average climate looks like, we can use the observed dataset and apply the changes to that – i.e. apply a 'change factor' or 'scaling factor'. This will create a dataset that represents a different climate – warmer or hotter, wetter or drier. A past change can be applied, or else a future projected change. If a particular threshold is important – for example an amount of annual rainfall or a particular average temperature for growing a crop, then this can be compared in the past, present and future climate datasets.

This simple example shows how we can take the annual average temperature of 1986-2005, then apply changes from the report (a 'scaling factor') to this data and see how this changes the area within the 25°C contour line. This can be relevant to real questions, for example the ideal growing conditions for Taro are above 25°C. All of Rarotonga is below 25°C in the annual average temperature in the current climate and the pre-industrial climate baseline of 1850-1900. Applying a scaling factor for projected change to a 2°C world, some of the coastal regions are higher than 25°C average annual temperature (the model average value of the projections is used: +1.3°C from 1850-1900 or +0.9°C from 1986-2005). We can also look at the range of possible change for 2050, from the lowest change projected for the very low emissions pathway (+0.3°C) to highest change projected under the very high emissions pathway (+2.3°C), shown in the bottom row of Figure 7.1.




Figure 7.1. The average annual temperature in 1986-2005 in Rarotonga (left: the 'WorldClim' high-resolution climate surface adjusted for the baseline used), and the area of the island above or below the 25°C boundary for various scenarios (right).

In addition to changes in the mean climate, impact assessment requires information about changes to extreme hazards, the exposure (e.g. people, assets, crops affected by the change in hazard) and vulnerability (the susceptibility to damage). For example, if projected changes in average temperature for 2050 are applied to historical daily temperature data, the output represents daily temperature data for 2050 which can be analysed for extreme events such as days over 30°C.

Hazard, exposure and vulnerability can be brought together to understand the impact, evaluate adaptation options and implement actions. This can be co-designed and investigated in a joint program with relevant stakeholders and researchers.

Box 3. Managing climate risks

According to the IPCC (2018), climate risks depend on the magnitude and rate of global warming, the exposure and vulnerability, and the implementation of adaptation and mitigation. Pacific Island nations are already dealing with climate risks, and these risks are projected to become much worse in future.

Limiting global warming to 1.5°C is projected to lower the risks compared to 2°C and beyond. Most adaptation needs will be lower for a global warming of 1.5°C. This would benefit marine, terrestrial, freshwater and coastal ecosystems, as well as human health, livelihoods, food security, water supply, security and economic growth. Pathways limited to 1.5°C global warming involve a decline in global net anthropogenic carbon dioxide emissions by about 45% from 2010 levels by 2030, reaching net zero around 2050.

A wide range of adaptation options is available. These can reduce the risks of climate change, but there are limits to adaptation and adaptive capacity for some human and natural systems. Strengthened multi-level governance, institutional capacity, policy instruments, technological innovation, transfer and mobilization of finance, and changes in human behaviour and lifestyles can enhance the feasibility of mitigation and adaptation.

Sustainable development supports fundamental societal and systems transitions and transformations that help limit global warming to 1.5°C. This can facilitate climate-resilient development pathways that achieve ambitious mitigation and adaptation outcomes. International cooperation is a critical enabler.



8 Data and Methods

For assessment of historical temperature in the Cook Islands region, we use five global observed datasets: HadCRUT5 (1850–2020; Morice et al., 2021), Berkeley Earth (1850–2019; Rohde and Hausfather, 2020), NOAA GlobalTemp (1880–2019; Huang et al., 2020; Zhang et al. 2020), Cowtan and Way (1850–2019; Cowtan and Way, 2014) and GISTEMP (1880–2019; Lenssen et al., 2019). For rainfall, we rely on the CMAP and GPCP merged gauge-satellite monthly precipitation datasets available from 1979 (see Yin et al. 2004 for a discussion and comparison of these), and ERA5 reanalysis rainfall data (1979–2020; Hersbach et al. 2020).

Gridded datasets use records from weather stations, satellite data and other sources, then fill in gaps in space and time. Therefore, the changes and trends in these gridded datasets agree with those in the underlying weather stations in general (see McGree et al. 2019) but are not exactly the same. Early periods include fewer weather observations with fewer supplementary data sources to draw upon, so these rely more heavily on the filling in across time and space and are therefore less reliable.

For the future climate, this report uses climate modelling output from the Coupled Model Inter-comparison Project phase 5 (CMIP5) of Taylor et al. (2012). The outputs from up to 36 climate models are used to show the ranges of possible change (Table 8.1), but detailed analyses and models we choose for scenario analyses are taken from a group of 21 models for RCP8.5 and RCP4.5, and 17 models for RCP2.6 that have passed evaluation tests (see Grose et al. 2014b), and we provide information about changes to climate features such as the SPCZ (Brown et al. 2013, 2020).

Projections are examined using three of the RCPs from van Vuuren et al., (2011): RCP2.6, RCP4.5 and RCP8.5 (there is also a RCP6.0, not used here). The RCPs are based on future pathways of all the major greenhouse gases, aerosols and land-use changes through the century, and are given a value based on the change in ‘radiative forcing’ by the year 2100 (Figure 8.1).




Figure 8.1. The enhanced greenhouse effect (or ‘radiative forcing’) in the three Representative Concentration Pathways (RCPs) used in this report. The three RCPs can be broadly thought of as a very high pathway with accelerating global emissions (RCP8.5), an intermediate pathway where emissions plateau (RCP4.5), and an ambitious decarbonisation pathway, roughly consistent with the Paris Agreement goal to keep global warming below 2°C (RCP2.6).

Average temperature and rainfall for the Cook Islands region are analysed and plotted in different ways to show historical and future context.

- To calculate changes at different 'global warming levels', e.g., 2°C global warming, 'time sampling' methods are employed (see James et al. 2017).
- Break points between different eras were detected by a statistical function that minimises the cost function (Matlab software: <https://au.mathworks.com/help/matlab/ref/ischange.html>).
- Representative climate scenarios for 2050 are derived from comparing the range of change in temperature and rainfall from the range of climate model outputs, and in understanding the most important drivers of change for this region, as described in the scientific literature. The dominant drivers of rainfall change are examined for the following regions:




Figure 8.2. Map of the western Pacific showing the sub-regions used for developing representative scenarios with attached storylines. Red line shows the average position of the centre of the SPCZ in the current climate, red dashed line shows the equator. Cook Islands South is in the “South of SPCZ” zone, outlined in light blue, and the Cook Islands North is in “On or north of the SPCZ”.

Table 8.1. CMIP5 models used in this report. While all models had RCP8.5 simulations available, a subset had RCP2.6 simulations, and the subset of models with evaluation information is also indicated.

No.	Name	Evaluated	RCP2.6	No.	Name	Evaluated	RCP2.6
1	'ACCESS1-0'	Yes		19	'GISS-E2-H'	Yes	Yes
2	'ACCESS1-3'	Yes		20	'GISS-E2-H-CC'		
3	'bcc-csm1-1'	Yes	Yes	21	'GISS-E2-R'	Yes	Yes
4	'bcc-csm1-1-m'		Yes	22	'GISS-E2-R-CC'		
5	'BNU-ESM'		Yes	23	'HadGEM2-AO'		Yes
6	'CanESM2'	Yes	Yes	24	'HadGEM2-CC'	Yes	
7	'CCSM4'	Yes	Yes	25	'HadGEM2-ES'	Yes	Yes
8	'CESM1-BGC'			26	'inmcm4'	Yes	
9	'CESM1-CAM5'		Yes	27	'IPSL-CM5A-LR'	Yes	Yes
10	'CMCC-CESM'			28	'IPSL-CM5A-MR'	Yes	Yes
11	'CMCC-CM'			29	'IPSL-CM5B-LR'		
12	'CMCC-CMS'			30	'MIROC5'	Yes	Yes
13	'CNRM-CM5'	Yes	Yes	31	'MPI-ESM-LR'	Yes	Yes
14	'FGOALS-s2'		Yes	32	'MPI-ESM-MR'		Yes
15	'FIO-ESM'		Yes	33	'MRI-CGCM3'	Yes	Yes
16	'GFDL-CM3'	Yes	Yes	34	'MRI-ESM1'		
17	'GFDL-ESM2G'	Yes	Yes	35	'NorESM1-M'	Yes	Yes
18	'GFDL-ESM2M'	Yes	Yes	36	'NorESM1-ME'	Yes	Yes

9 References

- Australian Bureau of Meteorology and CSIRO (2011). Climate Change in the Pacific: Scientific Assessment and New Research. Volume 1: Regional Overview. Volume 2: Country Reports
- Australian Bureau of Meteorology and CSIRO (2014). Climate Variability, Extremes and Change in the Western Tropical Pacific: New Science and Updated Country Reports. Pacific-Australia Climate Change Science and Adaptation Planning Program Technical Report, Australian Bureau of Meteorology and Commonwealth Scientific and Industrial Research Organisation, Melbourne, Australia.
- Bathiany, S., Scheffer, M., van Nes, E.H., Williamson, M.S. and Lenton, T.M. (2018). Abrupt Climate Change in an Oscillating World. *Scientific Reports* 8: 5040.
- Brown, J., Moise, A. and Colman, R. (2013). The South Pacific Convergence Zone in CMIP5 simulations of historical and future climate. *Climate Dynamics* 41: 2179-2197.
- Brown, J.R. Lengaigne, M., Lintner, B., Widlansky, M., van der Wiel, K., Duthel, C., Linsley, B., Matthews, A. and Renwick, J. (2020). South Pacific Convergence Zone dynamics, variability and impacts in a changing climate. *Nature Reviews Earth & Environment* 10.1038/s43017-020-0078-2.
- Chand S. S., Dowdy, A., Bell, S. and Tory, K. (2020). A Review of South Pacific Tropical Cyclones: Impacts of Natural Climate Variability and Climate Change. In: Lalit Kumar (eds) *Climate Change and Impacts in the Pacific*, Springer Climate. Springer, Cham. https://doi.org/10.1007/978-3-030-32878-8_6
- Cowtan, K. and Way, R.G. (2014). Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends. *Quarterly Journal of the Royal Meteorological Society*, 140(683), 1935–1944. <https://doi.org/10.1002/qj.2297>
- CSIRO and SPREP (2017). Climate change information for the Pacific: Guidance material to raise awareness and facilitate sectoral decision-making using science-based climate change information and services. Commonwealth Scientific and Industrial Research Organisation, Melbourne, Australia.
- CSIRO, Australian Bureau of Meteorology and SPREP (2015). Climate in the Pacific: a regional summary of new science and management tools, Pacific-Australia Climate Change Science and Adaptation Planning (PACCSAP) Program Summary Report. Melbourne, Australia, Commonwealth Scientific and Industrial Research Organisation.
- Frame, B., Lawrence, J., Ausseil, A.-G., Reisinger, A. and Daigneault, A. (2018). Adapting global shared socio-economic pathways for national and local scenarios. *Climate Risk Management* 21: 39-51.
- Frame, D., Joshi, M., Hawkins, E., Harrington, L.J. and de Roiste, M. (2017). Population-based emergence of unfamiliar climates. *Nature Climate Change* 7: 407-411.
- Grose, M.R., Bhend, J., Narsey, S., Gupta, A.S. and Brown, J.R. (2014a). Can We Constrain CMIP5 Rainfall Projections in the Tropical Pacific Based on Surface Warming Patterns? *Journal of Climate* 27: 9123-9138.
- Grose, M.R., Brown, J., Narsey, S., Brown, J., Murphy, B., Langlais, C., Sen Gupta, A., Moise, A. and Irving, D. (2014b). Assessment of the CMIP5 global climate model simulations of the western tropical Pacific climate system and comparison to CMIP3. *International Journal of Climatology* 34: 3382-3399.
- Grose, M.R., Narsey, S., Delage, F. P., Dowdy, A. J., Bador, M., Boschat, G., Chung, C., Kajtar, J. B., Rauniyar, S., Freund, M. B., Lyu, K., Rashid, H., Zhang, X., Wales, S., Trenham, C., Holbrook, N. J., Cowan, T., Alexander, L., Arblaster J. M. and Power S. (2020). Insights from CMIP6 for Australia's Future Climate. *Earth's Future* 8: e2019EF001469.
- Hawkins, E., (2018). Warming stripes. URL: <http://www.climate-lab-book.ac.uk/2018/warming-stripes>, 4(5).
- Hawkins, E., Frame, D., Harrington, L., Joshi, M., King, A., Rojas, M. and Sutton, R. (2020). Observed Emergence of the Climate Change Signal: From the Familiar to the Unknown. *Geophysical Research Letters* 47: e2019GL086259.
- Hersbach, H., Bell, W., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, Carole Peubey, J., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, R., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I.,

- Vamborg, F., Villaume, S. and Thépaut, J.-N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society* 146: 1999-2049.
- Huang, B., Menne, M.J., Boyer, T., Freeman, E., Gleason, B.E., Lawrimore, J.H., Liu, C., Rennie, J., Schreck, C., Sun, F., Vose, R., Williams, C., Yin, X., and Zhang, H. (2020). Uncertainty Estimates for Sea Surface Temperature and Land Surface Air Temperature in NOAA GlobalTemp Version 5. *Journal of Climate* 33: 1351-1379.
- IPCC (2013). Climate Change 2013: The Physical Science Basis. *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. T. F. Stocker, D. Qin, G.-K. Plattner et al. Cambridge, UK, and New York, NY, USA, Cambridge University Press.
- IPCC (2018). *Global warming of 1.5°C*. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)]. World Meteorological Organization, Geneva, Switzerland, 32 pp. www.ipcc.ch
- IPCC (2019). Special Report on Oceans and Cryosphere in a Changing Climate. www.ipcc.ch/srocc
- IPCC (2021). Climate Change 2021: The Physical Science Basis. *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. www.ipcc.ch
- James, R., Washington, R., Schleussner, C.-F., Rogelj, J. and Conway, D. (2017). Characterizing half-a-degree difference: a review of methods for identifying regional climate responses to global warming targets. *Wiley Interdisciplinary Reviews: Climate Change* 8: e457.
- King, A.D., Karoly, D.J. and Henley, B.J. (2017). Australian climate extremes at 1.5°C and 2°C of global warming. *Nature Climate Change* 7: 412.
- Knutson, T., Camargo, S.J., Chan, J.C., Emanuel, K., Ho, C.H., Kossin, J., Mohapatra, M., Satoh, M., Sugi, M., Walsh, K. and Wu, L. (2020). Tropical cyclones and climate change assessment: Part II. Projected response to anthropogenic warming. *Bulletin of the American Meteorological Society*, 101 (3): E303-E322.
- Lenssen, N.J.L., Schmidt, G.A., Hansen, J.E., Menne, M.J., Persin, A., Ruedy, R. and Zyss, D. (2019). Improvements in the GISTEMP Uncertainty Model. *Journal of Geophysical Research: Atmospheres*, 124(12), 6307–6326. <https://doi.org/10.1029/2018JD029522>
- McGree, S., Herold, N., Alexander, L., Schreider, S., Kuleshov, Y., Ene, E., Finaulahi, S., Inape, K., Mackenzie, B., Malala, H., Ngari, A., Prakash, B. and Tahani, L. (2019). Recent changes in mean and extreme temperature and precipitation in the western Pacific Islands. *Journal of Climate* 10.1175/JCLI-D-18-0748.1: JCLI-D-18-0748.0741.
- Menne, M.J., Williams, C.N., Gleason, B.E., Rennie, J.J., and Lawrimore, J.H. (2018). The Global Historical Climatology Network Monthly Temperature Dataset, Version 4. *Journal of Climate*, 31(24), 9835–9854. <https://doi.org/10.1175/JCLI-D-18-0094.1>
- Morice, C.P., Kennedy, J.J., Rayner, N.A., Winn, J.P., Hogan, E., Killick, R.E., Dunn, R.J.H., Osborn, T.J., Jones, P.D. and Simpson, I.R. (2021). An updated assessment of near-surface temperature change from 1850: the HadCRUT5 dataset. *Journal of Geophysical Research*. doi:10.1029/2019JD032361
- Rohde, R.A. and Hausfather, Z. (2020) The Berkeley Earth Land/Ocean Temperature Record. *Earth System Science Data* 12: 3469–3479.
- Shepherd, T.G., Boyd, E., Cabel, R., Chapman, S., Dessai, S., Dima-West, I., Fowler, H., James, R., Maraun, D., Martius, O., Senior, C., Sobel, A., Stainforth, D., Tett, S., Trenberth, K., van den Hurk, B., Watkins, N., Wilby, R. and Zenghelis, D. (2018). Storylines: an alternative approach to representing uncertainty in physical aspects of climate change. *Climatic Change* 151: 555-571.

Taylor, K.E., Stouffer, R.J. and Meehl, G.A. (2012). An Overview of CMIP5 and the Experiment Design. *Bulletin of the American Meteorological Society* 93: 485-498.


van Vuuren, D., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G., Kram, T., Krey, V., Lamarque, J., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. and Rose, S. (2011). The representative concentration pathways: an overview. *Climatic Change* 109: 5-31.

Yin, X.G., Gruber, A. and Arkin, P.A. (2004). Comparison of the GPCP and CMAP merged gauge-satellite monthly precipitation products for the period 1979-2001. *Journal of Hydrometeorology* 5: 1207-1222.

Zhang, H.-M., Huang, B., Lawrimore, J., Menne, M. and Smith, T.M. (2020). NOAA Global Surface Temperature Dataset (NOAAGlobalTemp), Version 5. NOAA National Centers for Environmental Information. doi:10.25921/9qth-2p70. Accessed May 2020.



© Saleasa Nihmei



www.rccap.org

www.pacificmet.net

www.pacificclimatechange.net

